arXiv:1303.1266v1 [quant-ph] 6 Mar 2013

Long-term effect of inter-mode transitions in quantum Markovian process

Sheng-Wen Li,[1] Li-Ping Yang,[2] and C. P. Sun[1]
[1]*Beijing Computational Science Research Center, Beijing 100084, China*
[2]*Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China*

We study the Markovian process of a multi-mode open system connecting with a non-equilibrium environment, which consists of several heat baths with different temperatures. As an illustration, we study the steady state of three linearly coupled harmonic oscillators in long time evolution, two of which contact with two independent bosonic heat baths with different temperatures respectively. We show that the inter-mode transitions mediated by the environment is responsible for the long time behaviour of the dynamics evolution, which is usually considered to take effect only in short time dynamics of the system immersed in a equilibrium heat bath with a single temperature. In non-equilibrium systems, the ignorance of the inter-mode transitions may give rise to some counter-intuitive result, thus they cannot be neglected.

PACS numbers: 03.65.Yz, 05.30.-d, 05.70.Ln

I. INTRODUCTION

When a small system contacts with a simple environment, i.e., a heat bath in canonical equilibrium with a temperature T, it would approach its canonical state with the same temperature T. This dynamics process, which erases the information of the initial state, is called *thermalization* [1–5]. The same thermalization process may also happen for a composite system, when all of its subsystems are coupled with the same common heat bath. The composite system would approach its canonical state of temperature T with respect to its total eigen modes.

However, nonequilibrium systems, e.g., the photosynthesis system, are more general in nature, and exhibit more rich physics [6–8]. A typical example is a composite system connecting with more than one heat baths with different temperatures. For a long-term evolution, the open composite system would not approach its canonical thermal state, but still it would be stabilized to a certain steady state. We call this process *non-thermal stabilization.*

In an open system, both equilibrium and non-equilibrium case as we mentioned above, the energy exchange with environment would mediate the transitions between the normal modes of the total system. It is usually believed that this inter-mode transition only takes effect on the dynamics of transient evolution within the time scale determined by the time-energy uncertainty [1, 9–12]. If this transition oscillates too fast in the relaxation time, averagely it has no contribution and can be neglected, which is known as rotating-wave approximation (RWA).

In this paper we consider whether or not such inter-mode transitions have a long-term effect on a composite open system contacting with a complicated environment, which consists of several independent heat baths with different temperatures. As an illustration, we study the steady state of three linearly coupled harmonic oscillators (HOs), two of which contact with two independent bosonic heat baths with different temperatures respectively. Specially, we are interested in the steady state of each local HO. It is found that if the inter-mode transition were ignored, there would be some counter-intuitive results in the long time steady state, especially when the frequency detuning and coupling strength of the local modes are small. In this regime, the coherent transitions among the total normal modes oscillate so slowly that they result in long-term effects, thus cannot be neglected.

The paper is arranged as follows. In Sec. II, we setup the model of the coupled system and give a master equation. In Sec. III, we show the stabilization result and make some analytical discussion by eliminating the degree of freedom of the mediating data bus. We propose a possible implementation in Sec. IV. The calculation is assisted by some properties of the characteristic description of Wigner function and Fokker-Planck equation. We leave these tricks in the appendices. Finally, summary is drawn in Sec. V.

II. MODEL SETUP

To study the long-term dynamics of a composite system coupled to a complicated environment, we use the coupled HOs system as an illustration. The system we study here is illustrated in Fig. 1. Two HO separate from each other for a distance. They are coupled indirectly by another mediating system, which is also modeled as a HO here. Effectively, we suppose the mediating HO does not contact with any environment. The three oscillators system can be described by a quadratic coupled Hamiltonian $H_S = H_0 + V$, where

$$\begin{aligned} H_0 &= \omega_L \hat{a}_L^\dagger \hat{a}_L + \omega_R \hat{a}_R^\dagger \hat{a}_R + \omega_{\mathrm{m}} \hat{b}^\dagger \hat{b}, \\ V &= g_L(\hat{a}_L^\dagger \hat{b} + \hat{a}_L \hat{b}^\dagger) + g_R(\hat{a}_R^\dagger \hat{b} + \hat{a}_R \hat{b}^\dagger), \end{aligned} \tag{1}$$

and H_0 describes the free Hamiltonian with local modes respectively defined by annihilation operators $\hat{a}_L$, $\hat{a}_R$ and $\hat{b}$; V describes the coupling among the local modes.

We assume the two oscillators locate remotely at different places, thus they may suffer from independent baths.



FIG. 1: (Color online) Demonstration of the coupled oscillators system. Two remotely located HOs are indirectly coupled by another mediating one. The HOs at the two ends contact with independent heat baths with different temperatures $T_{L/R}$.

We also assume each bath stays at a canonical thermal state with a temperature $T_{L/R}$. The whole system can be described by the total Hamiltonian $\mathcal{H} = H_S + H_B + V_{SB}$, where

$$\begin{aligned} H_B &= \sum_{\mathbf{k}_L} \omega_{\mathbf{k}_L} \hat{c}^\dagger_{\mathbf{k}_L} \hat{c}_{\mathbf{k}_L} + \sum_{\mathbf{k}_R} \omega_{\mathbf{k}_R} \hat{c}^\dagger_{\mathbf{k}_R} \hat{c}_{\mathbf{k}_R}, \\ V_{SB} &= \sum_{\sigma=L,R} \hat{a}^\dagger_\sigma \Gamma_\sigma + \hat{a}_\sigma \Gamma^\dagger_\sigma, \end{aligned} \tag{2}$$

and $\Gamma_\sigma = \sum_{\mathbf{k}_\sigma} g_{\mathbf{k}_\sigma} \hat{c}_{\mathbf{k}_\sigma}$. H_B is the free Hamiltonian of the two boson heat baths, each of which is modeled as a collection of boson modes, described by the boson annihilation operators $\hat{c}_{\mathbf{k}_\sigma}$. V_{SB} represents the linear coupling between the system and the environment.

We need to derive a master equation to study the dynamics of the open system. Actually for the coupled HO system, a correct treatment of the master equation should be based on the normal modes of H_S, but not the local modes $\hat{a}_{L/R}$ and $\hat{b}$. Otherwise, it may give rise to some counter-intuitive results. Thus, we diagonalize the Hamiltonian H_S as,

$$\begin{aligned} H_S &= (\hat{a}^\dagger_L,\, \hat{b}^\dagger,\, \hat{a}^\dagger_R) \begin{bmatrix} \omega_L & g_L & \\ g_L & \omega_{\mathrm{m}} & g_R \\ & g_R & \omega_R \end{bmatrix} \begin{pmatrix} \hat{a}_L \\ \hat{b} \\ \hat{a}_R \end{pmatrix} \\ &\equiv \mathbf{a}^\dagger \cdot \Omega \cdot \mathbf{a} = \sum_{i=1}^{3} \varepsilon_i \hat{A}^\dagger_i \hat{A}_i, \end{aligned} \tag{3}$$

where $\mathbf{a} = (\hat{a}_L,\, \hat{b},\, \hat{a}_R)^T$, and we also denote it as $(\hat{a}_1,\, \hat{a}_2,\, \hat{a}_3)^T$ hereafter (with redefined indices $1, 2, 3$ for $\hat{a}_L$, $\hat{b}$, $\hat{a}_R$ respectively). $\mathbf{A} = U \cdot \mathbf{a} = (\hat{A}_1,\, \hat{A}_2,\, \hat{A}_3)^T$ for A_i's being the normal modes. $U \cdot \Omega \cdot U^\dagger = \mathrm{diag}\,\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}$ gives the eigen frequencies of the normal modes. Although the normal modes are decoupled from each other in the isolated H_S, we can see below that the environment could mediately induce some effective couplings between these normal modes.

With the above notations, in Appendix A we derive a master equation to describe the long-term dynamics of the open system via Born-Markovian approximation [1]. In Schrödinger's picture, it reads as

$$\begin{aligned} \partial_t \rho =& i[\rho, \sum \varepsilon_i \hat{A}^\dagger_i \hat{A}_i] \\ &+ \sum_{ij} \frac{\Lambda^-_{ij}}{2} (2\hat{A}_i \rho \hat{A}^\dagger_j - \{\hat{A}^\dagger_j \hat{A}_i, \rho\}_+) \\ &+ \frac{\Lambda^+_{ij}}{2} (2\hat{A}^\dagger_i \rho \hat{A}_j - \{\hat{A}_j \hat{A}^\dagger_i, \rho\}_+), \end{aligned} \tag{4}$$

where

$$\begin{aligned} \Lambda^+_{ij} =& \frac{\gamma_L}{2} U_{i1} U^*_{j1} [N_L(\varepsilon_i) + N_L(\varepsilon_j)] \\ &+ \frac{\gamma_R}{2} U_{i3} U^*_{j3} [N_R(\varepsilon_i) + N_R(\varepsilon_j)], \\ \Lambda^-_{ji} =& \frac{\gamma_L}{2} U_{i1} U^*_{j1} [N_L(\varepsilon_i) + N_L(\varepsilon_j) + 2] \\ &+ \frac{\gamma_R}{2} U_{i3} U^*_{j3} [N_R(\varepsilon_i) + N_R(\varepsilon_j) + 2]. \end{aligned} \tag{5}$$

Here, $\gamma_\sigma(\varepsilon_i) = 2\pi J_\sigma(\varepsilon_i)$ characterizes the coupling strength with each bath, and $J_\sigma(\omega) = \sum_{\mathbf{k}_\sigma} |g_{\mathbf{k}_\sigma}|^2 \delta(\omega - \omega_{\mathbf{k}_\sigma})$ is the coupling distribution. For the usual case, we can assume that $\gamma_\sigma(\omega) \simeq \gamma_\sigma$ does not depend too much on ω and can be treated as constant. $N_\sigma(\omega) = [\exp(\omega/kT_\sigma) - 1]^{-1}$ is the Planck distribution for $\sigma = L, R$.

It is observed from the above master equation that the environment indeed induces an effective coupling between two normal modes A_i and A_j. $\Lambda^\pm_{ij}$ measure the transitions of the normal modes. This environment-mediating effect can be understood in the following way. The coupled system exchanges energy with environment through the interaction V_{SB}. Immediately after the normal mode A_i of the system emits an energy quanta ε_i to the environment, another process may happen in succession that the normal mode A_j absorbs back ε_j from the environment. Also, there is another possibility that mode A_i first absorb energy from the environment and then the energy is emitted through mode A_j. Thus, with the mediation of the environment, different normal modes A_i's of H_S are coupled. In fact, this effect of environment mediated coupling can be also found for two modes coupled to a common heat bath [13]. For comparison, we can also understand our result according to the case of common bath. Although there are two independent baths, the two baths are combined into a single one by the coupling between the local modes.

The transition terms with $i \neq j$ describes the effective coupling between different normal modes. In the interaction picture, these terms would contribute an oscillating factor $\exp[\pm i\delta\epsilon_{ij}\, t]$ resulted from the energy difference of the modes $\hat{A}_i$ and $\hat{A}_j$. This transition effect would be ignored if we apply RWA and drop these terms. However, as can be seen in the following, the energy difference $\delta\epsilon_{ij} \equiv \varepsilon_i - \varepsilon_j$ is determined by the coupling strengths and detunings of the local modes. When the interactions and detunings are small enough, $\delta\epsilon_{ij}$ becomes very small and the time dependent factors rotate so slowly that indeed these terms cannot be omitted.

Notice that $U_{in}\hat{a}_n = \hat{A}_i$, $U^*_{im}U_{in} = U_{mi}U^*_{ni} = \delta_{mn}$, we can also rewrite the master equation by the local modes $\hat{a}_i$,

$$\partial_t\rho = i[\rho,\, H_S] + \mathcal{L}_L[\rho] + \mathcal{L}_R[\rho], \tag{6}$$

where

$$\begin{aligned}\mathcal{L}_\sigma[\rho] =& \sum_n \frac{\tilde{\gamma}^+_{\sigma n}}{2}\big(\hat{a}^\dagger_\sigma\rho\hat{a}_n - \frac{1}{2}\{\hat{a}_n\hat{a}^\dagger_\sigma,\, \rho\}_+\big) + \mathbf{h.c.}\\ &+ \sum_n \frac{\tilde{\gamma}^-_{\sigma n}}{2}\big(\hat{a}_\sigma\rho\hat{a}^\dagger_n - \frac{1}{2}\{\hat{a}^\dagger_\sigma\hat{a}_n,\, \rho\}_+\big) + \mathbf{h.c.},\end{aligned}$$

and

$$\begin{aligned}\tilde{\gamma}^+_{\sigma n} =& \sum_i \frac{\gamma_\sigma}{2}U^*_{i\sigma}U_{in}\, N_\sigma(\varepsilon_i),\\ \tilde{\gamma}^-_{\sigma n} =& \sum_i \frac{\gamma_\sigma}{2}U_{i\sigma}U^*_{in}[N_\sigma(\varepsilon_i) + 1].\end{aligned}$$

Here σ takes L, R or $1, 3$. $\mathcal{L}_{L/R}[\rho]$ describes the effect from the left/right heat bath. For the local modes, besides the environment induced inter-mode coupling, the interaction between two modes can also give the inter-mode transition terms straightforwardly in the master equation. In this sense, it is concluded that the role of the environment mediation is to modify the direct inter-mode coupling as a kind of renormalization effect.

III. LONG-TERM STABILIZATION DYNAMICS

Comparing with the long time Markovian thermalization process in a heat bath with a single temperature, the present complicated environment with two temperatures $T_{L/R}$ cannot stabilize the whole system into a canonical thermal state. In this section, we first calculate the steady state of the open quantum system by straightforwardly solving the above master equation Eq. (4). Then we consider the mediating HO as a quantum data bus in the large detuning limit. The adiabatic elimination of this oscillator can formally induce a direct coupling between the left and right HOs. In this case, the analytical results about the stabilization can be obtained explicitly.

A. Steady state in long time limit

We now consider the indirect coupling case with a mediating data bus. The master equation without RWA can be solved with the help of the characteristic function of Wigner representation, which is defined as (see Appendix B),

$$\begin{aligned}\chi(\vec{\mu}) \equiv& \mathbf{Tr}\big[\rho\cdot\exp(\mathbf{A}^\dagger\cdot\vec{\mu} - \vec{\mu}^\dagger\cdot\mathbf{A})\big] \qquad (7)\\ =& \mathbf{Tr}\big[\rho\cdot\exp(\mathbf{a}^\dagger\cdot\vec{\kappa} - \vec{\kappa}^\dagger\cdot\mathbf{a})\big].\end{aligned}$$

Here, $\vec{\mu} = (\mu_1, \mu_2, \mu_3)^T$ and $\vec{\kappa} = (\kappa_1, \kappa_2, \kappa_3)^T$ are complex vectors with respect to the normal and local modes, and $\vec{\mu} = U\cdot\vec{\kappa}$. The corresponding Wigner function with three modes is defined as the Fourier transform of $\chi(\vec{\kappa})$,

$$W(\vec{\alpha}, \vec{\alpha}^*) = \frac{1}{(\pi^2)^3}\int d^2\vec{\kappa}\, e^{-\vec{\alpha}^\dagger\cdot\vec{\kappa} + \vec{\kappa}^\dagger\cdot\vec{\alpha}}\chi(\vec{\kappa}).$$

With this definition, we obtain the equation of $\chi(\vec{\mu})$ as

$$\partial_t\chi + \mathbf{z}\cdot\mathbf{T}\cdot\frac{\partial}{\partial\mathbf{z}^T}\chi = \mathbf{z}\cdot\mathbf{D}\cdot\mathbf{z}^T\chi, \tag{8}$$

where $\mathbf{z} = (\mu_1, \mu_2, \mu_3,\, \mu^*_1, \mu^*_2, \mu^*_3)$, and

$$\mathbf{T} = \begin{bmatrix} T^- & \mathbf{0}\\ \mathbf{0} & T^+\end{bmatrix}, \quad \mathbf{D} = \begin{bmatrix}\mathbf{0} & P\\ P^T & \mathbf{0}\end{bmatrix}.$$

$\mathbf{T}$ and $\mathbf{D}$ are 6×6 matrices with 3×3 blocks $T^\pm$ and P defined by

$$\begin{aligned}P_{ij} =& -\frac{1}{4}(\Lambda^-_{ij} + \Lambda^+_{ji}),\\ T^-_{ij} =& \frac{1}{2}(\Lambda^-_{ij} - \Lambda^+_{ji}) - i\varepsilon_i\delta_{ij},\\ T^+_{ij} =& \frac{1}{2}(\Lambda^-_{ji} - \Lambda^+_{ij}) + i\varepsilon_i\delta_{ij}.\end{aligned} \tag{9}$$

The equation (8) is the Fourier transformation of the Fokker-Planck equation about the Wigner function [14]. Formally we give the analytical solution for the steady state in Appendix C. Its expression is given as,

$$\chi(\vec{\mu}) = \chi(U\cdot\vec{\kappa}) = \exp\big[\mathbf{z}V^{-1}\cdot\mathbf{D}'\cdot(\mathbf{z}V^{-1})^T\big], \tag{10}$$

where V diagonalizes the matrix $\mathbf{T}$, i.e., $V\cdot\mathbf{T}\cdot V^{-1} = \mathrm{diag}\{\mathbf{d}_1, \cdots, \mathbf{d}_6\}$, and $\mathbf{D}'_{ij} = [V\mathbf{D}V^T]_{ij}/(\mathbf{d}_i + \mathbf{d}_j)$.

All the steady state properties of the composite system can be obtained from this formal solution Eq. (10). Specially, we are interested in the steady state of the HOs at the two ends. We can obtain $\chi_\sigma(\kappa_\sigma)$ for each local oscillator just by setting $\kappa_i = 0$ for all $i \neq \sigma$. Notice that V^{-1} and $\mathbf{D}'$ in the exponent of Eq. (10) are block diagonal and anti-diagonal respectively, thus it can be verified that $\chi_\sigma(\kappa_\sigma)$ is always of Gaussian form. Therefore, each oscillator finally stays at a state of the canonical distribution (see the proof in Appendix B),

$$\chi(\kappa_\sigma, \kappa^*_\sigma) = \exp\big[-(N^{\mathrm{eff}}_\sigma + \frac{1}{2})\,|\kappa_\sigma|^2\big]. \tag{11}$$

Since each HO can finally reach a steady state with a canonical form, we can treat it as an equivalent thermal state and define an effective temperature from its average occupation $N^{\mathrm{eff}}_\sigma = \langle\hat{a}^\dagger_\sigma\hat{a}_\sigma\rangle$,

$$T^{\mathrm{eff}}_\sigma = \omega_\sigma/\ln(1 + \frac{1}{N^{\mathrm{eff}}_\sigma}). \tag{12}$$

In Fig. 2(a, b) we show the effective temperatures of the oscillators at the two ends calculated from Eq. (10), changing with the detuning $\Delta = \omega_L - \omega_R$ and the coupling strengths $g_{L/R}$. Here we set $\overline{\omega} = (\omega_L + \omega_R)/2 \equiv 1$ as the energy scale and $\omega_{\mathrm{m}} = 2$.



FIG. 2: (Color online) The effective temperatures $T_L^{\rm eff}$ (lower blue ones) and $T_R^{\rm eff}$ (upper red ones) of the oscillators at the two ends calculated from the result without (a) (b) and with (c) RWA. We set $\varpi = (\omega_L + \omega_R)/2 = 1$ as the unit, and $T_L = 1$, $T_R = 3$, $\gamma_L = 0.002$, $\gamma_R = 0.003$, $\omega_{\rm m} = 2$. We set $g_L = g_R = g$ in (a) (c), and $g_L = g$, $g_R = 0.8g$ in (b). We plot four groups of curves according to $g = 0.02$, 0.04, 0.06, 0.08, distributed from outside to inside in these figures. At the small regime around the degeneracy point, RWA cannot give us a good enough result, especially when g is weak. The extremum points are shifted aside when $g_L \neq g_R$.

When the detuning Δ becomes large or when their coupling strength $g_{L/R}$ becomes small, the two oscillators tend to be thermalized with their own heat bath respectively. The effective temperatures get to the closest point around the resonance regime $\Delta \simeq 0$. This observation means that they are affected by the reservoir at the opposite side and heat transfer happens greatly. When $g_L = g_R$, the extremum points locate exactly at $\omega_L = \omega_R$, while they shift aside when $g_L \neq g_R$. When the interaction becomes strong, the effective temperatures of the two oscillators tend to get closer and closer, away from that of each heat bath.

As comparison, we also show some counter-intuitive observation resulting from the improper omission of the transition terms like $2\hat{A}_i\rho\hat{A}_j^\dagger - \{\hat{A}_j^\dagger\hat{A}_i, \rho\}$ with $i \neq j$ [see Fig. 2(c)]. In the large detuning area, this approximation shows well consistence with previous result in Fig. 2(a). But the effective temperatures always equals at the degeneracy point even when the coupling strength g is quite weak, i.e., when the oscillators tend to be decoupled from each other. Similar counter-intuitive result was also given in Ref. [5] for a coupled two-spin system.

With the above comparison, we look back at the master equation carefully, the transition terms like $e^{-i\delta\epsilon_{ij}t}\big(2\hat{A}_i\rho\hat{A}_j^\dagger - \{\hat{A}_j^\dagger\hat{A}_i, \rho\}_+\big)$ with $i \neq j$ contribute to the energy exchange of different modes $\hat{A}_i$ and $\hat{A}_j$, which can be characterized by $\langle\hat{A}_i^\dagger\hat{A}_j\rangle$. The oscillating factor at the front describes the phase of this transition. If the energy difference $\delta\epsilon_{ij} = \varepsilon_i - \varepsilon_j$ is large, this exchange oscillate so fast that this effect vanishes averagely. However, indeed $\delta\epsilon_{ij}$ is determined by the detuning and coupling strength. When Δ and $g_{L,R}$ are quite small, these terms cannot be omitted. Therefore, the transition between different normal modes plays an important role in such a composite non-equilibrium system with Markovian approximation. The rotating terms take effect to the long-term steady state behaviour and cannot be neglected, especially when the couplings between the subsystems are not quite strong.

It could be temporally concluded that people should be much careful to ignore the transition terms for a multi-mode open quantum system in a complicated environment. Otherwise, the correct physical effect may also be ignored.

B. Effective coupling in adiabatic limit

When the detuning of $\omega_{\rm m}$ to $\omega_{L/R}$ is large, we can apply Fröhlich-Nakajima transformation [15–17], and eliminate the mediating degree of freedom to get the following simplified Hamiltonian, which describes a system of two directly coupled oscillators,

$$H_S = \omega_L'\,\hat{a}_L^\dagger\hat{a}_L + \omega_R'\,\hat{a}_R^\dagger\hat{a}_R + g(\hat{a}_L^\dagger\hat{a}_R + \hat{a}_L\hat{a}_R^\dagger), \tag{13}$$

where

$$\begin{aligned}\omega_{L,R}' &= \omega_{L,R} + \frac{g_{L,R}^2}{\omega_{L,R} - \omega_{\rm m}}, \\ g &= \frac{1}{2}\Big(\frac{g_L g_R}{\omega_L - \omega_{\rm m}} + \frac{g_L g_R}{\omega_R - \omega_{\rm m}}\Big).\end{aligned} \tag{14}$$

We can see that the coupling strengths g_L and g_R contribute a correction to the renormalized frequencies $\omega_{L,R}'$. For this simplified Hamiltonian, we can write down the analytical expression of eigen frequencies ε_i and the transformation U for the normal modes $\hat{A}_\pm$. Denoting $\omega_L' = \varpi - \Delta/2$, $\omega_R' = \varpi + \Delta/2$, we have

$$\begin{aligned}&\varepsilon_\pm = \varpi \pm \tilde{\Delta}_g, \qquad \tilde{\Delta}_g = \Big(\frac{\Delta^2}{4} + g^2\Big)^{\frac{1}{2}}, \\ &U = \begin{bmatrix}\alpha & \beta \\ \beta & -\alpha\end{bmatrix}, \quad \frac{2\alpha\beta}{\alpha^2 - \beta^2} = \frac{2g}{\Delta}.\end{aligned} \tag{15}$$

It follows from Eq. (15) that the energy difference $\delta\epsilon_{ij} = 2\tilde{\Delta}_g$ depends on the detuning Δ and coupling strength g. When Δ and g are small, the factors $\exp[\pm 2i\tilde{\Delta}_g t]$ of the transition terms between the two normal modes oscillate quite slowly. Therefore, these transition terms may not be omitted.

We carry out the similar calculation for this simplified two oscillators system as previously, which gives an explicit expression for the steady state of each oscillator, described by the characteristic function $\chi_\sigma(\kappa_\sigma)$,

$$\begin{aligned}\chi_\sigma(\kappa_\sigma) =& \exp\big[-(N_\sigma^{\mathrm{eff}}+\frac{1}{2})\,|\kappa_\sigma|^2\big], \qquad (16)\\ N_\sigma^{\mathrm{eff}} =& \big[\mathsf{A}_\sigma N_L(\varepsilon_-)+\mathsf{B}_\sigma N_L(\varepsilon_+)\\ &+\mathsf{C}_\sigma N_R(\varepsilon_-)+\mathsf{D}_\sigma N_R(\varepsilon_+)\big]/\Phi.\end{aligned}$$

Here N_σ^{eff} is the occupation number of the effective thermal distribution ($\sigma = L, R$) determined by the linear combination of $N_{L,R}(\varepsilon_\pm)$, and the coefficients are,

$$\Phi = \gamma_L\gamma_R(\gamma_L+\gamma_R)^2+16\tilde{\Delta}_g^2(\alpha^2\gamma_L+\beta^2\gamma_R)(\beta^2\gamma_L+\alpha^2\gamma_R),$$

$$\begin{aligned}\mathsf{A}_L &= \alpha^2\big[\gamma_L\gamma_R(\gamma_L+\gamma_R)^2+16\alpha^2\tilde{\Delta}_g^2(\beta^2\gamma_L+\alpha^2\gamma_R)\gamma_L\big],\\ \mathsf{B}_L &= \beta^2\big[\gamma_L\gamma_R(\gamma_L+\gamma_R)^2+16\beta^2\tilde{\Delta}_g^2(\alpha^2\gamma_L+\beta^2\gamma_R)\gamma_L\big],\\ \mathsf{C}_L &= 16\alpha^2\beta^2\tilde{\Delta}_g^2(\beta^2\gamma_L+\alpha^2\gamma_R)\gamma_R,\\ \mathsf{D}_L &= 16\alpha^2\beta^2\tilde{\Delta}_g^2(\alpha^2\gamma_L+\beta^2\gamma_R)\gamma_R,\end{aligned}$$

and

$$\begin{aligned}\mathsf{A}_R &= 16\alpha^2\beta^2\tilde{\Delta}_g^2(\beta^2\gamma_L+\alpha^2\gamma_R)\gamma_L,\\ \mathsf{B}_R &= 16\alpha^2\beta^2\tilde{\Delta}_g^2(\alpha^2\gamma_L+\beta^2\gamma_R)\gamma_L,\\ \mathsf{C}_R &= \beta^2\big[\gamma_L\gamma_R(\gamma_L+\gamma_R)^2+16\beta^2\tilde{\Delta}_g^2(\alpha^2\gamma_L+\beta^2\gamma_R)\gamma_R\big],\\ \mathsf{D}_R &= \alpha^2\big[\gamma_L\gamma_R(\gamma_L+\gamma_R)^2+16\alpha^2\tilde{\Delta}_g^2(\beta^2\gamma_L+\alpha^2\gamma_R)\gamma_R\big].\end{aligned}$$

From Eq. (16) we see that each oscillator achieves a canonical state. Especially, at the degeneracy point $\omega_L' = \omega_R'$, we have $\alpha^2 = \beta^2 = 1/2$, and the difference of the populations is,

$$\begin{aligned}& N_L^{\mathrm{eff}} - N_R^{\mathrm{eff}}\\ &= \frac{\big[N_L(\varepsilon_+)-N_R(\varepsilon_+)\big]+\big[N_L(\varepsilon_-)-N_R(\varepsilon_-)\big]}{2(1+4g^2/\gamma_L\gamma_R)}. \qquad (17)\end{aligned}$$

The above equation (17) obviously shows that the effective temperatures of the two oscillators are not equal at the degeneracy point.

This analytical result Eq. (16) gives the same profile with Fig. 2(a, b). From the above arguments, we clearly see that the effective temperature of each HO is determined by the competition between the interaction of the oscillators and that with their reservoirs, which is characterized by their ratio $g^2/\gamma_L\gamma_R$. When the interaction strength g of the oscillators is large, the difference of the temperatures tend to be zero, i.e., the interaction between the oscillators tend to bind them as a whole to be stabilized, and the two oscillators tend to get equal temperatures. Moreover, since $\omega_{L,R}'$ is renormalized by the interaction energy $g_{L,R}$, at the degeneracy point $\omega_L = \omega_R$, we may have $\omega_L' \neq \omega_R'$ if $g_L \neq g_R$ [see Eq. (14)]. Therefore, the extremum points are shifted aside from $\omega_L = \omega_R$ as shown in Fig. 2(b).



FIG. 3: (Color online) Two remotely located NAMRs are indirectly coupled via a superconducting TLR. The bias voltage provides a difference between the NAMR and the TLR, so they can be coupled by a capacitance. At the same time, the voltage noise provides each NAMR with independent heat bath which has different effective temperature.

From the discussion above, we conclude that the inter-mode transitions have long-term effect on the composite system in a complicated nonequilibrium environment. This is quite different from some previous viewpoint that the inter-mode transitions only have transient effect within the time scale determined by the time-energy uncertainty, which usually applies in conventional thermalization process [1, 9–12].

IV. REMARKS AND CONCLUSION

Before we summarize the stabilization result of the coupled oscillator system in a non-equilibrium environment, we discuss a possible implementation by two nanomechanical oscillators (NAMR) connected via a superconducting transmission line (TLR) to test the theoretical results here, as shown in Fig. 3.

The electromagnetic field inside the TLR may be treated as several boson modes whose frequencies are discretely distributed [18, 19]. Only one of the TLR modes, which is nearly resonant to the NAMRs, can couple with the NAMRs effectively. For example, the voltage distribution of the lowest even mode along the TLR is,

$$V(x) = \sqrt{\frac{\omega_{\mathrm{m}}}{cL}}\cos\frac{2\pi x}{L}\,(\hat{b}+\hat{b}^\dagger), \qquad (18)$$

where $\omega_{\mathrm{m}} = 2\pi/L\sqrt{lc}$ is the frequency of this mode, L is the length of the TLR, and l, c are the inductance and capacitance per unit length. The voltage gets maximum at the two ends, where the NAMRs are coupled with the TLR via a displacement dependent capacitance [20–24]. The vibration mode of each NAMR may also be treated as a single boson. To the lowest order, C_x depends linearly on the movement of the NAMR, $C_x \simeq C_x^0(1+\hat{x}/d_0)$. Applying a voltage bias V_g to the NAMRs, we have the interaction as

$$H_{\mathrm{int}} = \frac{1}{2}C_x^0(1+\frac{\hat{x}}{d_0})\big(V(x)-V_g\big)^2. \qquad (19)$$

Quantizing the coordinate of the NAMR as $\hat{x} = \delta x_0(\hat{a}+$

$\hat{a}^\dagger$), we obtain an interaction term as $H_{\text{int}} = g(\hat{a}+\hat{a}^\dagger)(\hat{b}+\hat{b}^\dagger)$.

The quality factor of the TLR is usually much higher than that of the NAMRs, thus we neglect the dissipation of the TLR. The voltage noises on the two sides, which arise from resistance and bring in the Joule heat, may provide the NAMRs with independent heat baths with different effective temperatures , and this can be controlled and measured in experiment [25–27]. Typically, the frequencies of the NAMRs and TLR are around hundreds of MHz to several GHz, depending on the sizes of the devices, and the coupling strengths are usually several MHz [18, 19, 28]. In this regime, it is appropriate to make Jaynes-Cummings approximation to obtain linear coupling as the above analysis.

In summary, we have studied the long-term behaviour of a coupled HO system connecting with a complicated environment which consists of two independent heat baths with different temperatures. We deduced a master equation with respect to the normal modes. With the help of the characteristic description of Wigner function, we obtained the numerical and analytical results for the steady state of each local oscillator. These results show that much attention must be payed when we apply RWA in a nonequilibrium system. When the coupling of the subsystems is weak, the inter-mode transitions belong the RWA may have long-term effect to the steady state thus cannot be neglected. This is different from the case in conventional thermalization problems, where only one canonical heat bath is involved. The non-thermal stabilization process is determined by the competition between the rate of the inter-mode transition and that of the energy exchange with each private heat bath.

This work is supported by National Natural Science Foundation of China under Grants Nos.11121403, 10935010 and 11074261, and National 973-program Grants No. 2012CB922104.

Appendix A: Derivation of master equation

We show the derivation of the master equation Eq. (4) here. In the interaction picture of H_S+H_B, the interaction with the environment becomes,

$$\begin{aligned} V_I(t) &= V_I^L(t)+V_I^R(t), \qquad \text{(A1)}\\ V_I^\sigma(t) &= \hat{a}_\sigma^\dagger(t)\Gamma_\sigma(t)+\hat{a}_\sigma(t)\Gamma_\sigma^\dagger(t), \end{aligned}$$

where

$$\begin{aligned} \hat{a}_\sigma(t) &= \sum_j U_{j\sigma}^* \hat{A}_j e^{-i\varepsilon_j t}, \qquad \text{(A2)}\\ \Gamma_\sigma(t) &= \sum_{\mathbf{k}_\sigma} g_{\mathbf{k}_\sigma}\hat{c}_{\mathbf{k}_\sigma} e^{-i\omega_{\mathbf{k}_\sigma}t}. \end{aligned}$$

Here $\hat{A}_i = U_{ij}\hat{a}_j$ are the normal modes of the interacting oscillators system.

We take Born-Markovian approximation and put these interaction terms into the following equation [1],

$$\begin{aligned} \partial_t\rho =& -\int_0^\infty d\tau\,\mathbf{Tr}_B\left[V_I(t),[V_I(t-\tau),\rho(t)\otimes\rho_B]\right] \qquad \text{(A3)}\\ =& -\int_0^\infty d\tau\,\mathbf{Tr}_B\left[V_I^L(t),\left[V_I^L(t-\tau),\rho(t)\otimes\rho_B\right]\right]\\ & -\int_0^\infty d\tau\,\mathbf{Tr}_B\left[V_I^R(t),\left[V_I^R(t-\tau),\rho(t)\otimes\rho_B\right]\right]. \end{aligned}$$

Here we assume that the state of each bath is a canonical thermal one and does not change with time, i.e., $\rho_B = \rho_B^L\otimes\rho_B^R$, $\rho_B^\sigma \propto \exp[-H_B^\sigma/kT_\sigma]$ and $H_B^\sigma = \omega_{\mathbf{k}_\sigma}\hat{c}_{\mathbf{k}_\sigma}^\dagger\hat{c}_{\mathbf{k}_\sigma}$ where $T_{L/R}$ is the temperature of the left/right heat bath. Thus, terms like $\mathbf{Tr}_B[V_I^L(t)V_I^R(t-\tau)\rho(t)\otimes\rho_B]$ always vanish, because they only contain the first moment of each bath.

The rightside of Eq. (A3) contains two integrals of the same form. Each integral gives four terms, one of which is calculated bellow as an example,

$$\begin{aligned} &\int_0^\infty d\tau\,\mathbf{Tr}_B\left[\hat{a}_\sigma^\dagger(t)\Gamma_\sigma(t)\cdot\rho(t)\otimes\rho_B\cdot\hat{a}_\sigma(t-\tau)\Gamma_\sigma^\dagger(t-\tau)\right]\\ =&\int_0^\infty d\tau\,\hat{a}_\sigma^\dagger(t)\rho(t)\hat{a}_\sigma(t-\tau)\left\langle\Gamma_\sigma^\dagger(t-\tau)\Gamma_\sigma(t)\right\rangle_B, \qquad \text{(A4)} \end{aligned}$$

where

$$\begin{aligned} \hat{a}_\sigma^\dagger(t)\rho(t)\hat{a}_\sigma(t-\tau) &= \sum_{i,j} U_{i\sigma}U_{j\sigma}^*\,\hat{A}_i^\dagger\rho\hat{A}_j\,e^{i(\varepsilon_i-\varepsilon_j)t}\cdot e^{i\varepsilon_j\tau},\\ \left\langle\Gamma_\sigma^\dagger(t-\tau)\Gamma_\sigma(t)\right\rangle_B &= \sum_{\mathbf{k}_\sigma}\left|g_{\mathbf{k}_\sigma}\right|^2\langle\hat{c}_{\mathbf{k}_\sigma}^\dagger\hat{c}_{\mathbf{k}_\sigma}\rangle_{\text{th}}\,e^{-i\omega_{\mathbf{k}_\sigma}\tau} \qquad \text{(A5)}\\ &= \int_0^\infty d\omega\,J_\sigma(\omega)N_\sigma(\omega)\,e^{-i\omega\tau}. \end{aligned}$$

Here, $J_\sigma(\omega) = \sum_{\mathbf{k}_\sigma}\left|g_{\mathbf{k}_\sigma}\right|^2\delta(\omega-\omega_{\mathbf{k}_\sigma})$ is the coupling spectrum, and $N_\sigma(\varepsilon_i) = [\exp(\varepsilon_i/kT_\sigma)-1]^{-1}$ is the Planck distribution with temperature T_σ. The integral Eq. (A4) gives

$$\begin{aligned} &\sum_{i,j} U_{i\sigma}U_{j\sigma}^*\,\hat{A}_i^\dagger\rho\hat{A}_j\,e^{i(\varepsilon_i-\varepsilon_j)t}\\ &\quad\times\frac{\gamma_\sigma(\varepsilon_j)}{2}\cdot N(\varepsilon_j)+i\mathbf{P}\int_0^\infty d\omega\,\frac{J_\sigma(\omega)N_\sigma(\omega)}{\varepsilon_j-\omega}. \qquad \text{(A6)} \end{aligned}$$

Here, we denote $\gamma_\sigma(\varepsilon_i) = 2\pi J_\sigma(\varepsilon_i)$, which characterizes the coupling strength with each heat bath. The principle integral contributes to Lamb shift, and we omit this term in this paper.

The physical meaning of Eq. (A4) may be understood in the following way. At time $t-\tau$, the coupled HO system emits energy to the environment, and then absorbs back at time t. However, the energy exchange with the environment during this process is done by the total normal modes $\hat{A}_i$ but not the local modes $\hat{a}_i$. Thus, when the emission and absorption modes are not the same one, there is an oscillating factor $\exp[i(\varepsilon_i - \varepsilon_j)t]$ left. $\delta\epsilon_{ij} \equiv \varepsilon_i - \varepsilon_j$ characterizes the splitting amplitude resulting from the coupling. By the mediation of the environment, the different normal modes $\hat{A}_i$ of the system are coupled together.

Other terms of Eq. (A3) can be also obtained as above. Each of the two integrals gives the following Lindblad-like form with an extra oscillating factor,

$$\begin{aligned}
&\sum_{ij} \frac{1}{2} U_{i\sigma} U^*_{j\sigma} \Big(\gamma_\sigma(\varepsilon_i)[N_\sigma(\varepsilon_i)+1] + \gamma_\sigma(\varepsilon_j)[N_\sigma(\varepsilon_j)+1]\Big) \\
&\qquad \times \Big(\hat{A}_j \rho \hat{A}_i^\dagger - \frac{1}{2}\{\hat{A}_i^\dagger \hat{A}_j,\, \rho\}_+\Big) e^{i\delta\epsilon_{ij} t} \\
&+ \sum_{ij} \frac{1}{2} U_{i\sigma} U^*_{j\sigma} [\gamma_\sigma(\varepsilon_j)\, N_\sigma(\varepsilon_i) + \gamma_\sigma(\varepsilon_j)\, N_\sigma(\varepsilon_j)] \qquad \text{(A7)} \\
&\qquad \times \Big(\hat{A}_i^\dagger \rho \hat{A}_j - \frac{1}{2}\{\hat{A}_j \hat{A}_i^\dagger,\, \rho\}_+\Big) e^{i\delta\epsilon_{ij} t}.
\end{aligned}$$

For simplicity, we assume $\gamma_\sigma(\varepsilon_i) \simeq \gamma_\sigma$ does not depend too much on ω and can be treated as constant. In sum of Eqs. (A3, A7), we get the following master equation in Schrödinger's picture, and the oscillating factors do not appear,

$$\begin{aligned}
\partial_t \rho =& i[\rho, \sum \varepsilon_i \hat{A}_i^\dagger \hat{A}_i] \\
&+ \sum_{ij} \frac{\Lambda^-_{ij}}{2} \big(2\hat{A}_i \rho \hat{A}_j^\dagger - \{\hat{A}_j^\dagger \hat{A}_i, \rho\}_+\big) \qquad \text{(A8)} \\
&+ \frac{\Lambda^+_{ij}}{2} \big(2\hat{A}_i^\dagger \rho \hat{A}_j - \{\hat{A}_j \hat{A}_i^\dagger, \rho\}_+\big),
\end{aligned}$$

where

$$\begin{aligned}
\Lambda^+_{ij} =& \frac{\gamma_L}{2} U_{i1} U^*_{j1} [N_L(\varepsilon_i) + N_L(\varepsilon_j)] \\
&+ \frac{\gamma_R}{2} U_{i3} U^*_{j3} [N_R(\varepsilon_i) + N_R(\varepsilon_j)], \qquad \text{(A9)} \\
\Lambda^-_{ji} =& \frac{\gamma_L}{2} U_{i1} U^*_{j1} [N_L(\varepsilon_i) + N_L(\varepsilon_j) + 2] \\
&+ \frac{\gamma_R}{2} U_{i3} U^*_{j3} [N_R(\varepsilon_i) + N_R(\varepsilon_j) + 2].
\end{aligned}$$

The terms with $i \neq j$ describes the transition between different normal modes. These terms are often omitted by RWA.

Appendix B: Characteristic function of Wigner representation

The Wigner representation often give us great convenience to study properties of quantum oscillators. It can be defined from a characteristic function [29],

$$\chi_w(\kappa, \kappa^*) = \mathbf{Tr}\big[e^{\kappa \hat{a}^\dagger - \kappa^* \hat{a}} \rho\big] \qquad \text{(B1)}$$

The Wigner function is defined as the Fourier transform of $\chi_w(\kappa, \kappa^*)$,

$$W(\alpha, \alpha^*) = \frac{1}{\pi^2} \int d^2\kappa\, e^{-\kappa\alpha^* + \kappa^*\alpha} \chi_w(\kappa, \kappa^*). \qquad \text{(B2)}$$

For a system that consists of two oscillators, the characteristic function can be defined as,

$$\chi_{12}(\kappa_1, \kappa_2) = \mathbf{Tr}_{12}\big[e^{\kappa_1 \hat{a}_1^\dagger - \kappa_1^* \hat{a}_1} \cdot e^{\kappa_2 \hat{a}_2^\dagger - \kappa_2^* \hat{a}_2} \rho_{12}\big].$$

From this definition, immediately we can find that if we had known $\chi_{12}(\kappa_1, \kappa_2)$ for the whole system explicitly, it would be quite easy to get the description of the subsystems $\chi_{1(2)}$, just by setting $\kappa_{2(1)} = 0$ in $\chi_{12}(\kappa_1, \kappa_2)$, without having to calculate the reduced density matrix of subsystems $\rho_{1(2)}$. This provides a simple method for us to study the state of subsystems.

Besides, for the thermal state of a oscillator $\rho_T = \mathcal{Z}^{-1} \exp[-\frac{\omega}{kT} \hat{a}^\dagger \hat{a}]$, the characteristic function is,

$$\chi_T(\kappa, \kappa^*) = \exp\big[-(N + \frac{1}{2})\, |\kappa|^2\big]. \qquad \text{(B3)}$$

Here $N = [\exp(\omega/kT) - 1]^{-1}$ is the Planck distribution.

As seen from the definition, $\chi_w(\kappa, \kappa^*)$ and $W(\alpha, \alpha^*)$ can be mapped into each other through Fourier transformation. It is also well known that there is one-to-one correspondence between a physical Wigner function and a density matrix. Therefore, there is one and only one density matrix ρ decided by a legal $\chi_w(\kappa, \kappa^*)$.

Thus, if we have a characteristic function $\chi(\kappa, \kappa^*)$ which has a Gaussian form like Eq.(B3), with $N \geq 0$, we can always come into the fact that the corresponding density matrix is

$$\rho = \frac{1}{\mathcal{Z}} \sum_{n=0}^{\infty} e^{-n\,\beta^{\text{eff}}\omega} |n\rangle\langle n|, \qquad \text{(B4)}$$

where β^{eff} comes from $N = \big[\exp(\beta^{\text{eff}}\omega) - 1\big]^{-1}$. This is a canonical state for the oscillator with an effective temperature $1/\beta^{\text{eff}}$. A more rigorous proof lies bellow.

Proof: From the definition of $\chi_w(\kappa, \kappa^*)$, we have

$$\begin{aligned}
\chi_w(\kappa, \kappa^*) &= e^{\frac{1}{2}|\kappa|^2} \mathbf{Tr}\big[e^{\kappa \hat{a}^\dagger} \rho e^{-\kappa^* \hat{a}}\big] \qquad \text{(B5)} \\
&= e^{\frac{1}{2}|\kappa|^2} \int \frac{d^2\alpha}{\pi} e^{\kappa\alpha^* - \kappa^*\alpha} \langle\alpha|\rho|\alpha\rangle.
\end{aligned}$$

If we have a Gaussian formed characteristic function like Eq.(B3), we can correspondingly get $\langle\alpha|\rho_T|\alpha\rangle$ by reversed transformation of the equation above,

$$\begin{aligned}
\langle\alpha|\rho_T|\alpha\rangle &= \int \frac{d^2\kappa}{\pi} e^{-\kappa\alpha^* + \kappa^*\alpha} \cdot \exp\big[-(N+1)\,|\kappa|^2\big] \\
&= \frac{1}{N+1} \exp\big[-\frac{|\alpha|^2}{N+1}\big]. \qquad \text{(B6)}
\end{aligned}$$

On the other hand, we can also expand $\langle\alpha|\rho_T|\alpha\rangle$ as,

$$\begin{aligned}
\langle\alpha|\rho_T|\alpha\rangle &= \sum_{m,n} \langle\alpha|m\rangle \,\langle m\,|\rho_T|\,n\rangle\, \langle n|\alpha\rangle \qquad \text{(B7)} \\
&= \sum_{m,n} \langle m\,|\rho_T|\,n\rangle \cdot \frac{(\alpha^*)^m \alpha^n}{\sqrt{m!n!}} e^{-|\alpha|^2}.
\end{aligned}$$

Comparing with the expansion of Eq. (B6), we can get the matrix elements of ρ_T,

$$\langle m\,|\rho_T|\,n\rangle=\frac{(1+\frac{1}{N})^{-n}}{N+1}\delta_{mn}. \tag{B8}$$

Denote $\exp[\beta^{\text{eff}}\omega]=1+\frac{1}{N}$, we can see that ρ_T is a canonical state. ■

Appendix C: Steady solution of Fokker-Planck equation

The standard form of Fokker-Planck equation and its characteristic equation are as follows,

$$\frac{\partial P}{\partial t}+\sum_i\lambda_i\frac{\partial}{\partial y_i}(y_iP)=\frac{1}{2}\sum_{ij}\sigma_{ij}\frac{\partial^2P}{\partial y_i\partial y_j}, \tag{C1}$$

$$\frac{\partial f}{\partial t}-\sum_i\lambda_i\xi_i\frac{\partial f}{\partial\xi_i}=-\frac{1}{2}f\sum_{ij}\sigma_{ij}\xi_i\xi_j,\quad \mathbf{Re}\lambda_i<0.$$

$f(\vec{\xi},t)$ is the Fourier transformation of $P(\vec{y},t)$,

$$f(\vec{\xi},t)=\int d^ny\,P(\vec{y},t)e^{-i\vec{\xi}\cdot\vec{y}}.$$

The equation of $f(\vec{\xi},t)$ is a first-order quasi-linear partial differential one. It can be solved analytically [30], and the solution is,

$$f(\vec{\xi},t)=\Phi(\xi_ie^{\lambda_it},\cdots)\cdot\exp\Big[-\frac{1}{2}\sum_{ij}\sigma_{ij}\frac{\xi_i\xi_j}{\lambda_i+\lambda_j}\Big]. \tag{C2}$$

$\Phi(\cdots)$ is determined according to the initial condition, and $\Phi(t\to\infty)=1$.

In our problem, the equation of the characteristic function is

$$\partial_t\chi+\vec{z}\cdot\mathbf{T}\cdot\frac{\partial}{\partial\vec{z}^T}\chi=\vec{z}\cdot\mathbf{D}\cdot\vec{z}^T\chi. \tag{C3}$$

The only difference with the standard form is that $\mathbf{T}$ is not diagonal here. We first diagonalize it and make it a standard form. Denoting $V\cdot\mathbf{T}\cdot V^{-1}=\text{diag}\{\mathbf{d}_1,\cdots,\mathbf{d}_n\}$ and $\vec{z}=\vec{\xi}\cdot V$, we can transform our equation into the standard Fokker-Planck form,

$$\partial_t\chi+\vec{\xi}\cdot\mathbf{d}\cdot\frac{\partial}{\partial\vec{\xi}^T}\chi=\vec{\xi}\cdot V\mathbf{D}V^T\cdot\vec{\xi}^T\chi. \tag{C4}$$

Now we could write down the steady solution as

$$\chi(\vec{z})=\exp\left[\vec{z}V^{-1}\cdot\mathbf{D}'\cdot(\vec{z}V^{-1})^T\right], \tag{C5}$$

$$\text{where}\qquad \mathbf{D}'_{ij}=\frac{[V\mathbf{D}V^T]_{ij}}{\mathbf{d}_i+\mathbf{d}_j}.$$

[1] H. Breuer and F. Petruccione, *The theory of open quantum systems* (Oxford University Press, 2002).

[2] M. Rigol, V. Dunjko, and M. Olshanii, Nature **452**, 854 (2008).

[3] N. Linden, S. Popescu, A. J. Short, and A. Winter, Phys. Rev. E **79**, 061103 (2009).

[4] J.-Q. Liao, H. Dong, and C. P. Sun, Phys. Rev. A **81**, 052121 (2010).

[5] J.-Q. Liao, J.-F. Huang, and L.-M. Kuang, Phys. Rev. A **83**, 052110 (2011).

[6] F. Caruso, A. W. Chin, A. Datta, S. F. Huelga, and M. B. Plenio, J. Chem. Phys. **131**, 105106 (2009).

[7] S. Yang, D. Z. Xu, Z. Song, and C. P. Sun, J. Chem. Phys. **132**, 234501 (2010).

[8] J.-Q. Liao, J.-F. Huang, L.-M. Kuang, and C. P. Sun, Phys. Rev. A **82**, 052109 (2010).

[9] J. Jing, Z.-G. Lü, and Z. Ficek, Phys. Rev. A **79**, 044305 (2009).

[10] Z.-H. Li, D.-W. Wang, H. Zheng, S.-Y. Zhu, and M. S. Zubairy, Phys. Rev. A **80**, 023801 (2009).

[11] Q. Ai, Y. Li, H. Zheng, and C. P. Sun, Phys. Rev. A **81**, 042116 (2010).

[12] Y. Li, J. Evers, H. Zheng, and S.-Y. Zhu, Phys. Rev. A **85**, 053830 (2012).

[13] D. P. S. McCutcheon, A. Nazir, S. Bose, and A. J. Fisher, Phys. Rev. A **80**, 022337 (2009).

[14] D. F. Walls and G. J. Milburn, *Quantum optics*, 2nd ed. (Springer, 2008).

[15] H. Fröhlich, Phys. Rev. **79**, 845 (1950).

[16] S. Nakajima, Adv. Phys. **4**, 363 (1955).

[17] Z.-L. Xiang, X.-Y. Lu, T.-F. Li, J. Q. You, and F. Nori, arXiv:1211.1827 (2012).

[18] A. Blais, R.-S. Huang, A. Wallraff, S. M. Girvin, and R. J. Schoelkopf, Phys. Rev. A **69**, 062320 (2004).

[19] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R.-S. Huang, J. Majer, S. Kumar, S. M. Girvin, and R. J. Schoelkopf, Nature **431**, 162 (2004).

[20] M. R. Geller and A. N. Cleland, Phys. Rev. A **71**, 032311 (2005).

[21] C. P. Sun, L. F. Wei, Y.-X. Liu, and F. Nori, Phys. Rev. A **73**, 022318 (2006).

[22] L. F. Wei, Y.-X. Liu, C. P. Sun, and F. Nori, Phys. Rev. Lett. **97**, 237201 (2006).

[23] L. Tian, M. S. Allman, and R. W. Simmonds, New J. Phys. **10**, 115001 (2008).

[24] L. Tian, Phys. Rev. B **79**, 193407 (2009).

[25] F. Giazotto, T. T. Heikkilä, A. Luukanen, A. M. Savin, and J. P. Pekola, Rev. Mod. Phys. **78**, 217 (2006).

[26] Y.-X. Chen and S.-W. Li, Europhys. Lett. **97**, 40003 (2012).

[27] F. Giazotto and M. J. Martínez-Pérez, Nature **492**, 401 (2012).

[28] Y. Greenberg, Y. A. Pashkin, and E. Il'ichev, Phys. Uspekhi **55**, 382 (2012).

[29] C. Gardiner and P. Zoller, *Quantum noise*, Vol. 56 (Springer, 2004).

[30] M. C. Wang and G. E. Uhlenbeck, Rev. Mod. Phys. **17**, 323 (1945).